

# Effectiveness of Water Extract of Sea Lada (*Ulva Lactuca*) from Aceh Waters to Reduce Blood Glucosa Levels in Diabetic Rats

<sup>1</sup>Rinawati, <sup>2</sup>Sri Wahyuni Muhsin, <sup>3</sup>Hanif Muchdatul Ayunda

<sup>1</sup>Nutrition Department, Public Health Faculty, Universitas Teuku Umar, Indonesia, [rinawati@utu.ac.id](mailto:rinawati@utu.ac.id)

<sup>2</sup>Nutrition Department, Public Health Faculty, Universitas Teuku Umar, Indonesia, [sriwahyunimuhsin@utu.ac.id](mailto:sriwahyunimuhsin@utu.ac.id)

<sup>3</sup>Nutrition Department, Public Health Faculty, Universitas Teuku Umar, Indonesia, [hanifmuchdatul@utu.ac.id](mailto:hanifmuchdatul@utu.ac.id)

**Corresponding author:** Rinawati, e-mail: [rinawati@utu.ac.id](mailto:rinawati@utu.ac.id)

## ABSTRACT

Diabetes mellitus (DM) is a non-communicable disease characterized by high blood glucose levels caused by damage to the pancreas, which cannot produce insulin. Antidiabetic drugs and insulin injections in the long term have specific side effects, so herbal medicine is considered safer to use. Sea lettuce (*Ulva lactuca*) is one of the herbal medicines that have the potential as an antidiabetic. Methods: Experimental using a completely randomized design. Rats were divided into five groups with three replicates, namely: Kn = normal control, Ka = STZ control, P1 = DM rats + 200 mg/KgBB extract dose, P2 = DM rats + 300 mg/KgBB extract dose, and P3 = DM rats + 400 mg/KgBB extract dose. The stages of this study began with the preparation of tools and materials, phytochemical tests, the preparation of test animals and testing extracts on test animals, and the measurement of blood glucose levels in rats before and after they were given extracts. Blood glucose levels were measured using a glucometer. Data analysis was performed statistically (ANOVA). The results of the phytochemical tests detected were steroids, saponins, and phenols/tannins, while alkaloids, terpenoids, and flavonoids were not. The most effective blood glucose level measurement result is at a dose of 400 mg/KgBB

## ARTICLE INFORMATION

**Submitted:** 08/11/2022

**Revised:** 20/11/2022

**Accepted:** 27/11/2022

**Published Online:** 28/11/2022

**Keywords:**

Diabetic

*Ulva lactuca*

Water extract

**How to cite this article:** Rinawati, R., Muhsin, S. W., & Ayunda, H. M. (2022). Effectiveness of Water Extract of Sea Lada (*Ulva Lactuca*) from Aceh Waters to Reduce Blood Glucosa Levels in Diabetic Rats. *Journal of Nutrition Science*, 3(2), 60-66. doi:10.35308/jns.v3v2.6833

## Introduction

Diabetes mellitus is a non-communicable disease caused by damage to the pancreas so that it cannot synthesize insulin needed for glucose breakdown. Impaired insulin production will affect blood glucose levels that continue to increase and are not controlled. Although not contagious, diabetes is one of the most dangerous diseases that cause death. Diabetes can cause various disease complications, such as eye damage, vascular disorders, heart disease, kidney failure, and, most fatally, as a cause of death (Thongsom, 2013). The prevalence of diabetes, mainly types 2 diabetes, has increased globally by 30%. The most common complication experienced by people with diabetes is acute heart failure. (Busui, 2022). Hyperglycemia also affects the performance of glands such as the hypothalamus, pituitary gland, gonads, and perigonads. Glands that cannot function properly cause a decrease in the secretion of gonadal steroids such as GnRH, FSH, LH, and testosterone, thus affecting male fertility and reproductive health

(Andlip, 2023). Hyperglycemia is the leading cause of various vascular complications through the following mechanisms: activation of the polyol and hexosamine pathways; activation of protein kinase C; increased oxidative stress; increased production of advanced glycation end products; increased synthesis of growth factors, cytokines, and angiotensin II. In addition, diabetic complications are also due to the involvement of epigenetic mechanisms and micro RNAs in pathogenesis (Chiarelli, 2023).

According to the FDI (International Diabetes Federation), people with diabetes aged 20–79 years amounted to approximately 366 million in 2011 and are expected to rise to 522 million people in 2030 (Whiting, DR. 2011). Data compiled by the WHO (2018) shows that there were 2.2 million cases of death caused by diabetes in 2012, making it the seventh leading cause of death worldwide. The age range of people with diabetes is below 70 years. It is estimated that there will be an increase in the

number of diabetics by 25% in 2030 and 51% in 2045 (Saeedi, 2019).

Diabetes treatment is still relatively expensive and sometimes has side effects, so herbal plants are starting to be tried as alternative drugs because they have hypoglycemic properties and fewer side effects. Herbal plants are spread throughout Asia but are most abundant in Indonesia, which has 80% of the world's population. There are 2000-7500 species of herbal plants estimated to be distributed in Indonesia. Most people use herbal plants in the form of herbal drinks or herbal medicine. In addition, herbal plants are also used as raw materials for drugs that have been standardized after undergoing research and clinical trials in humans (Zahra, H., 2022). People use herbal plants as antidiabetics by boiling the leaves with water to make herbal medicine (Rinawati, 2019). Researchers have done much research on medicinal plants because their bioactive content gives them biological effects so that they can be used as antioxidants, antibacterials, antidiabetics, anticancer agents, etc. One of the many research projects conducted by researchers today is the potential of herbal plants as antidiabetics (Prisdiani, Y. 2021).

WHO (World Health Organization) states that treatment using natural ingredients has been used by 60% of the population of developed countries (DepKes, RI, 2007). One of the natural ingredients that can potentially be medicinal is sea lettuce (*Ulva lactuca*). *Ulva lactuca* is algae found in marine waters and is usually processed into food ingredients by the community. *Ulva lactuca* is one type that is classified as "Father Seaweed" or a type of green algae that can be consumed and has antioxidant, antifungal, antibacterial, and antitumor content (Arbi et al., 2016). Because it has a high carbohydrate and fat content but a low-fat content, *Ulva lactuca* has the potential as a functional food raw material. Pigments contained in *Ulva lactuca* can also act as antioxidants that inhibit cancer, obesity, and diabetes; these pigments include neoxanthin, antheraxanthin, and vaucheriaxanthin, which have allenic bonds like fucoxanthin (Costa, 2018).

According to some studies, *Ulva lactuca* can be used as an antihyperglycemic in DM disease. The methanol extract of *Ulva lactuca* is proven to reduce hyperglycemia in vitro (Murugesan, S. 2015), and the ethanol extract of *Ulva lactuca* can reduce cholesterol levels in DM rats (Fairuz, 2014) and help restore body weight in diabetic rats who have lost weight (Rinawati, 2020). *Ulva lactuca* can reduce blood glucose levels by inhibiting the post-prandial glucose absorption process due to its bioactive content's activity (Ditasari, 2022). The oligosaccharide content of *Ulva lactuca* effectively restores blood glucose metabolic processes. It delays brain aging in elderly patients with type 2 DM through the GLP-1/GLP-1R pathway to mobilize communication between the brain and gut (Chen, 2022). *Ulva lactuca* ethanol extract contains alkaloids, steroids, and phenolics/tannins (Zulfadhli,

2018). Research on *Ulva lactuca* using an aqueous extract against hyperglycemia and its phytochemical screening has yet to be carried out, especially in the Aceh area.

## Method

### Place of Research

his research was conducted at the Laboratory of the Academy of Pharmacy, YPPM Mandiri Banda Aceh, for testing the water extract of *Ulva lactuca* on test rats *in vivo*.

### Tools and Materials

The tools used in this study are sonde needles, syringes, cages, weight scales to measure the development of rat body weight during and after treatment, a blood glucose level checker brand called "Easy Touch," and Easy Touch brand blood glucose level measuring strips, distilled water, male rats with an average weight of 200 grams, rat food, and STZ to condition rats into diabetes.

## Research Stages

### Preparation of sea lettuce extract

After being cleaned of dirt attached to the leaves, *Ulva lactuca* is thoroughly washed and drained. The drying process of *Ulva lactuca* is done by air drying it for several days until it is completely dry and brittle. The powder is made by crushing it in a blender and then sifting it to get a powder that is not too coarse or too fine.

### A Phytochemical Test of Sea Lettuce

Water-solvent maceration method for extracting *Ulva lactuca* the juice solution was then evaporated with a vacuum rotary evaporator at 40 °C until a thick extract was obtained. Phytochemical testing was carried out using the Harborne method (1997).

### Animal Test Preparation

First, rats were given nicotinamide (NA) solution at a dose of 230 mg/kgBB. After 15 minutes, rats were injected with streptozotocin (STZ) at a dose of 65 mg/kg BB. The STZ required to induce depends on the rats' body weight (BW).

### *In vivo* testing of extracts on test animals

*Ulva lactuca* water extract was given to test animals *in vivo* using a sonde needle at doses of 200 mg/kg, 300 mg/kg, and 400 mg/kg for 30 days. Every week blood glucose levels were checked every five days.

## Research design

This research is experimental with a completely randomized design. The total number of test animals was 25, divided into five treatments with three replicates:

Kn = Normal control (only given food and distilled water).

Ka = STZ control (rats induced by 65 mg STZ)

P1 = DM rats + 200 mg/KgBB extract dose

P2 = DM rats plus a 300 mg/KgBB extract dose.

P3 = DM rats + 400 mg/KgBB extract dose

## Parameters observed

### 1. Blood glucose levels before and after STZ induction

Measuring blood glucose levels before and after STZ induction aims to determine whether STZ has worked perfectly to cause hyperglycemia. Usually, mice will experience an increase in blood glucose on the third day after STZ induction (Saputra, 2028).

### 2. Glucose levels before and after administration of the extract.

Measuring glucose levels before and after administration of the extract aims to determine whether the administration of *Ulva lactuca* water extract effectively reduces blood glucose levels in DM rats.

## Data Analysis

Data analysis was done descriptively in the form of tables and graphs. Data processing and analysis using computer aid Anova analysis were performed for the measurement of blood glucose levels in each group; if there were differences between treatment groups, they were further analyzed by the DMRT (Duncan multiple range test).

## Results

### Phytochemical Content of Sea Lettuce

Sea lettuce has a thin, sheet-like structure resembling transparent green plastic. In making extracts, sea lettuce must first be cleaned of adhering dirt and then air-dried for several days. After drying, sea lettuce is crushed into powder using a blender and then ready for extraction. The extraction process on sea lettuce is done using water as a solvent. The phytochemical content contained in sea lettuce water extract is as follows:

Table 1: Phytochemical test results of sea lettuce (*Ulva lactuca*) water extract.

Metabolite Content	Reagen	Test Result	Observation Results
Alkaloid	Mayer	-	No white precipitate formed
	Wagner Dragendorff	-	No brown precipitate formed. No red precipitate formed
Steroid	Liebermann-Burchard	+	Green Color Formed
	Test	-	
Terpenoid	Liebermann-Burchard	-	There was no red color formed
	Test	-	
Saponin	Aquades	+	Foaming
Flavonoid	HCl and Mg Metal	-	No Formation Reddish Color
	Test	-	
Phenol/Tannin	FeCl3	+	Green Color Formed

Based on the table above, the phytochemical content of sea lettuce water extract is not all detected; of the six metabolite contents that are checked, only three are positive, namely steroids, saponins, and phenols/tannins while alkaloids, terpenoids, and flavonoids are not detected. These results are similar to previous research conducted by Zulfadhli (2018) on the extraction of sea lettuce using ethanol solvents, where the results of phytochemical screening contain alkaloid, steroid, and tannin compounds. Water solvents are polar and

are one of the solvents allowed because they meet pharmaceutical-grade requirements and are safe to use (Latifa, 2022). The polarity of the water solvent is because it has a hydroxyl group (Buhang, 2019). Alkaloids are semipolar (Hesturini, 2022), flavonoids are polar (Suryandari, 2022), and terpenoids are nonpolar, so they will not dissolve in polar solvents (Resti, 2022). Although flavonoids are polar, they were not detected in this study. The amount is tiny in *Ulva lactuca*, so water solvents cannot draw it.

## Blood Glucose Levels

In this study, rats were divided into the Normal Control group, STZ Control, P1, P2, and P3. The normal control group is a rat test animal only given aquadest without any treatment. This aims to compare the difference between animals that receive treatment and animals that do not. STZ control is a test animal that is always conditioned diabetes as a comparison of the decrease in blood glucose levels in rats treated with *Ulva lactuca* water extract. Group P1 is a diabetic rat given a dose of 200 mg / kgBB, Group P2 is a diabetic rat given a dose of 300 mg / kgBB, and P3 is a diabetic rat group given a dose of 400 mg / kgBB.

The measurement of blood glucose levels starts from day 0 to day 28. Blood sampling is done in 7 days. Measurement of day 0 is a measurement of blood glucose levels carried out after rats are acclimatized and have not received any treatment. Measurement on day 7 is the measurement of blood glucose levels carried out after the test animals were induced by STZ (SZ Control Group, P1, P2, and P3). Day 14, 21, and 28, measurements of glucose levels were taken when the test animals were already in a diabetic state. The results of blood glucose level measurements are presented in the following table:

Table 2: Blood Glucose Level Measurement Results

Treatment	Rat Glucose Level (mg/dl)				
	Day 0	7	14	21	28
Normal control (KN)	80,33±5,03 <sup>d</sup>	80,33±2,51 <sup>a</sup>	79,33±4,04 <sup>d</sup>	79,33±4,04 <sup>a</sup>	80,67±0,57 <sup>a</sup>
STZ control (Ka)	92,33±4,93 <sup>c</sup>	247,33±2,51 <sup>d</sup>	251,00±1,73 <sup>c</sup>	262,00±2,64 <sup>b</sup>	271,00±7,93 <sup>a</sup>
Dose 200 mg (P1)	110,33±4,50 <sup>b</sup>	290,00±5,00 <sup>b</sup>	281,67±2,88 <sup>a</sup>	274,67±4,50 <sup>a</sup>	245,00±5,00 <sup>b</sup>
Dose 300 mg (P2)	106,67±3,51 <sup>a</sup>	301,00±3,60 <sup>a</sup>	268,67±3,51 <sup>b</sup>	230,00±4,35 <sup>c</sup>	216,67±6,50 <sup>c</sup>
Dose 400 mg (P3)	81,33±1,15 <sup>d</sup>	277,67±3,21 <sup>c</sup>	245,00±8,66 <sup>c</sup>	207,00±8,71 <sup>d</sup>	107,00±7,00 <sup>d</sup>

Notes: Numbers in the same row followed by the same superscript letter are not significantly different at the 5% test level.

## Discussion

The results of measuring blood glucose levels in rat test animals on day 0 are shown in the table above and show no significant difference between treatments. This is because the rats in each treatment are still in a normal state and have not been given any treatment. Although there are differences in glucose levels in each group, they are still normal. The different physiological states of each rat can cause the difference.

The increase in blood glucose levels in STZ-induced rats occurred when measuring day 7, while the

blood glucose levels of the KN group remained normal. This is because the rats in the KN treatment group were conditioned to be normal and were only used as a comparison. The increase in blood glucose levels in the Ka, P1, and P2 groups is due to the induction of STZ, which has begun to damage the pancreas of rats so that it disrupts insulin secretion, which ultimately causes glucose levels in the blood of rats to increase above normal limits.

The results of measuring blood glucose levels in rat test animals on day 0 are shown in the table above and show no significant difference between treatments. This is because the rats in each treatment are still in a normal state and have not been given any treatment. Although there are differences in glucose levels in each group, they are still normal. The different physiological states of each rat can cause the difference.

The increase in blood glucose levels in STZ-induced rats occurred when measuring day 7, while the blood glucose levels of the KN group remained normal. This is because the rats in the KN treatment group were conditioned to be normal and were only used as a comparison. The increase in blood glucose levels in the Ka, P1, and P2 groups is due to the induction of STZ, which has begun to damage the pancreas of rats so that it disrupts insulin secretion, which ultimately causes glucose levels in the blood of rats to increase above normal limits.

If the glucose level in the test animal exceeds the normal limit, this indicates that the animal is diabetic. According to Port (1995), normal blood glucose levels are 110 mg/dL when fasting or 140 mg/dL 2 hours after eating, whereas diabetes occurs when glucose levels exceed 126 mg/dL when fasting or 200 mg/dL when not fasting. STZ induction can trigger oxidative stress that ultimately disrupts pancreatic performance and causes high glucose levels (hyperglycemia). There are three mechanisms of oxidative stress, as follows: (1) non-enzymatic protein glycation; (2) aldose reductase pathway, which uses the SDH (sorbitol dehydrogenase) enzyme to convert sorbitol to fructose. Sorbitol degradation is slow, so sorbitol accumulates in cells. This causes an increase in osmotic pressure, which ultimately damages cells (3). The change in oxidative status is characterized by changes in endogenous antioxidant activity and increased oxidative damage to biomolecules. To reduce oxidative damage, antioxidants are needed (Setiawan, 2005).

An observation of the effectiveness of *Ulva lactuca* water extract was done after administration of the extract on the 8th day. On day 14, blood glucose levels in the Ka, P1, P2, and P3 groups remained high, but the Ka and P3 groups were not significantly different from the other groups. This data shows that *Ulva lactuca* water extract in the first week has been unable to reduce blood glucose levels to normal in the DM rat group. e normal limit, this indicates that the animal is diabetic. According to Port (1995), normal blood glucose levels are 110

mg/dL when fasting or 140 mg/dL 2 hours after eating, whereas diabetes occurs when glucose levels exceed 126 mg/dL when fasting or 200 mg/dL when not fasting. STZ induction can trigger oxidative stress that ultimately disrupts pancreatic performance and causes high glucose levels (hyperglycemia). There are three mechanisms of oxidative stress, as follows: (1) non-enzymatic protein glycation; (2) aldose reductase pathway, which uses the SDH (sorbitol dehydrogenase) enzyme to convert sorbitol to fructose. Sorbitol degradation is slow, so sorbitol accumulates in cells. This causes an increase in osmotic pressure, which ultimately damages cells (3). The change in oxidative status is characterized by changes in endogenous antioxidant activity and increased oxidative damage to biomolecules. To reduce oxidative damage, antioxidants are needed (Setiawan, 2005).

An observation of the effectiveness of *Ulva lactuca* water extract was done after administration of the extract on the 8th day. On day 14, blood glucose levels in the Ka, P1, P2, and P3 groups remained high, but the Ka and P3 groups were not significantly different from the other groups. This data shows that *Ulva lactuca* water extract in the first week has been unable to reduce blood glucose levels to normal in the DM rat group.

The measurement of blood glucose levels at week 21 showed that each treatment group differed significantly. The Ka group experienced an increase in blood glucose levels. Although blood glucose levels in groups P1, P2, and P3 are still classified as hyperglycemia, there is a decrease compared to day 14 measurements.

The measurement of glucose levels on day 28 also showed a significant difference. Group Ka still experienced an increase in glucose levels from 262,00±2,64 mg/kgBB on day 21 to 271,00±7,93. The P1 and P2 groups experienced a decrease in blood glucose levels, but the levels were still high. The P3 group has reached normal limits but is slightly higher than the KN group. The percentage reduction in blood glucose levels can be calculated through the difference between the measurement results on day 7 and day 28. The data is presented in the following table:

Table 3: Percentage reduction in blood glucose levels of diabetic rats

Group	Decrease in Glucose Blood Glucose Level (%)
Normal Control (KN)	0%
STZ Control (Ka)	Increase 10%
Dose 200 mg (P1)	16%
Dose 300 mg (P2)	28%
Dose 400 mg (P3)	61%

Based on the table above, the highest percentage decrease in blood glucose levels occurred in the P3 group at a dose of 400 mg/kg BW, as much as 61%, followed by the P2 group by 28%. The lowest

percentage decrease was in the P1 group, which was only able to reduce blood glucose levels by 16% after being given *Ulva lactuca* water extract for 21 days. KN did not experience any changes because it was used as a treatment control, while Ka, a diabetic control, experienced an increased blood glucose level of 10%. This is because the rats were diabetic and were not given any treatment. The decrease in blood glucose levels in the group treated with *Ulva lactuca* water extract can be caused by the content of phytochemical compounds that can help regenerate pancreatic cells. According to Tair (2018), *Ulva lactuca* has hypotensive, hypoglycemic, and antiatherogenic properties that can improve or prevent cardiovascular complications because it contains polysaccharides. The polysaccharide content can play a role in improving the antioxidant pathway of diabetic rats that experience complications of neuropathy syndrome (Appu, A., 2022).

The results of phytochemical screening showed that *Ulva lactuca* water extract was positive for steroids, saponins, phenols, and tannins. There may be other phytochemicals besides those mentioned earlier. However, they most likely cannot be attracted by water solvents, or their content is so small that it is not detected during phytochemical screening. Steroids, saponins, and phenol/tanine are antioxidant compounds that can counteract free radicals to repair damage to pancreatic cells to a better or even normal state. The regenerated pancreas causes insulin production to increase so that it can reduce hyperglycemia (Dyahnugra, 2015).

In diabetes, saponins have several functions, including (a) helping improve blood glucose absorption, reducing blood glucose and serum insulin levels, and improving glucose tolerance. (b) help restore weight loss due to diabetes, improve insulin resistance, and help increase the proportion of protein kinase. (c) modulate lipid metabolism, promote adipocyte thermogenesis, restore insulin sensitivity and glucose homeostasis, and reduce inflammation. (d) aid increases insulin secretion. (e) has anti-inflammatory properties and inhibits platelet aggregation. (f) has an activity to reduce plasma glucose and total cholesterol levels (Guzman, D.C., 2020). Saponins can also reduce oxidative stress levels and regulate insulin signaling pathways (Jiang, 2020).

Tannins have hypoglycemic activity by increasing the glycogenesis process. Tannins also function as astringents (chelators), which can wrinkle the epithelial membrane of the small intestine, thereby reducing the absorption of food juice reduced sugar absorption results in a slower rate of blood sugar rise (Suryani, 2013). Furthermore, tannins lower blood glucose levels by delaying intestinal glucose absorption by inhibiting  $\alpha$ -amylase and  $\alpha$ -glucosidase activity in insulin-sensitive tissues (Tandi, J., 2020). The mechanism of steroid use to reduce blood glucose levels is by helping the

pancreas produce and secrete insulin (Munawaroh, 2022). Steroids also act as antioxidants and overcome pancreatic beta-cell damage, the main cause of insulin non-production (Novalinda, 2022). Secondary metabolites in *Ulva lactuca* water extract work together to cause hypoglycemic effects (Pasaribu, 2012).

### Conclusion

The chemical compounds contained in the water extract of sea lettuce (*Ulva lactuca*) are steroids, saponins, and phenols/tannins. A water extract of sea lettuce (*Ulva lactuca*) can reduce glucose levels in diabetic rats, with the most effective dose being 400 mg/kgBB. Further tests must be done with different dose ranges to get a more effective dose. Besides that, it is necessary to add a normal control group given *Ulva lactuca* water extract treatment to see if it affects weight loss and blood glucose levels.

### Acknowledgement

The Beginner Lecturer Research Grant funded the author's research from DRP2M DIKTI in 2018–2019.

### References

- Agus, Rheza Paleva. "Mekanisme Resistensi Insulin Terkait Obesitas." *Jurnal Ilmiah Kesehatan Sandi Husada* 8.2 (2019): 354-358.
- Andlib, Nida, et al. "Abnormalities in sex hormones and sexual dysfunction in males with diabetes mellitus: A mechanistic insight." *Acta Histochemica* 125.1 (2023): 151974.
- Appu, Anoop, et al. "Injectable chitosan-*Ulva lactuca* polysaccharide hydrogel neutralizes oxidative stress in experimental diabetic neuropathy." *GSC Biological and Pharmaceutical Sciences* 21.1 (2022): 042-059.
- Arbi, Basyrowi, Widodo Farid Ma'ruf, and Romadhon Romadhon. "AKTIVITAS SENYAWA BIOAKTIF SELADA LAUT (*Ulva lactuca*) SEBAGAI ANTIOKSIDAN PADA MINYAK IKAN" *Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology* Vol 12 No 1. 2016: 12-18.
- Chen, Y., e. al, "Regulatory Mechanisms Of The Green Alga *Ulva Lactuca* Oligosaccharide Via The Metabolomics And Gut Microbiome In Diabetic Mice". *Current Research In Food Science*, Volume 5, 14 July 2022: 1127-1139.
- Chen, Yihan, et al. "Regulatory mechanisms of the green alga *Ulva lactuca* oligosaccharide via the metabolomics and gut microbiome in diabetic mice." *Current Research in Food Science* 5 (2022): 1127-1139.
- Da Costa, Junet Franzisca, Windu Merdekawati, and Ferly Rambu Otu. "Analisis Proksimat, Aktivitas Antioksidan, Dan Komposisi Pigmen *Ulva Lactuca* L. Dari Perairan Pantai Kukup." *Jurnal Teknologi Pangan dan Gizi* 17.1 (2018): 1-17.

- Departemen Kesehatan Republik Indonesia. 2007. Keputusan Menteri Kesehatan Republik Indonesia Nomor 381 Tahun 2007 tentang Kebijakan Obat Tradisional Nasional Tahun 2007. Jakarta: Depkes RI.
- Ditasari, Ulvi, Agung Giri Samudra, and Reza Pertiwi. "EFEKTIVITAS ANTIHIPERGLIKEMIA EKSTRAK ETANOL *Ulva* sp. DAN *Sargassum* sp. PADA MENCIT YANG DIINDUKSI SUKROSA: ANTIHYPERGLYCHEMIC EFFECTIVENESS OF ETHANOL EXTRACT *Ulva* sp. AND *Sargassum* sp. IN SUCROSE-INDUCED MICULES." *Medical Sains: Jurnal Ilmiah Kefarmasian* 7.4 (2022): 789-796.
- Dyahnugra, A.A dan Widjarnato, S.B. "Pemberian Ekstrak Bubuk *Simplisia* kulit manggis (*Garcinia Mangostana* L) Menurunkan Kadar Glukosa Pada Tikus Putih (*Rattus norvegicus*) Strain Wistar Jantan Kondisi Hiperglikemik". Jurnal publikasi. Volume 3, No 1. 2015: 122
- Guzmán, David Calderón, et al. "Consumption of cooked common beans or saponins could reduce the risk of diabetic complications." *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy* 13 (2020): 3481.
- Hesturini, Rosa Juwita, et al. "ANALGESIC TEST AND TOXICITY OF n-HEXANA FRACTION TREMBESI LEAVES (*Samanea saman* (Jacq.) Merr.) IN MICE (*Mus musculus* L.)." *Jurnal Farmasi Sains dan Praktis* (2022): 32-41.
- Jiang, Shuang, et al. "Antidiabetic effect of *Momordica charantia* saponins in rats induced by high-fat diet combined with STZ." *Electronic Journal of Biotechnology* 43 (2020): 41-47.
- Latifa, Nabila Nur, Lanny Mulqie, and Siti Hazar. "Penetapan Kadar Sari Larut Air Dan Kadar Sari Larut Etanol *Simplisia* Buah Tin (*Ficus carica* L.)." *Bandung Conference Series: Pharmacy*. Vol. 2. No. 2. 2022.
- Munawwaroh, Siti Winda, Sri Peni Fitrianiingsih, and Ratu Choesrina. "Studi Literatur Aktivitas Antidiabetes Biji Mahoni (*Swietenia mahagoni* (L.) Jacq.)." *Bandung Conference Series: Pharmacy*. Vol. 2. No. 2. 2022.
- Murugesan, S., Anand, B.M., Bhuvanewari, S., Kotteswari, M., andThennarasan, S."In vitro Antidiabetic Activity of Methanolic Extracts of Selected Marine Algae". *European Journal of Pharmaceutical and Medical Research*. Volume 2, No.6. 2015: 256-260.
- Nilsson, Peter M., and Andrea Natali. "Insulin and Blood Pressure Relationships." *Blood Pressure Disorders in Diabetes Mellitus*. Springer, Cham, 2023. 119-128.
- Novalinda, Novalinda, Mukti Priastomo, and Laode Rijai. "Literature Review: Bahan Alam yang Berpotensi sebagai Antidiabetes: Literature Review: Natural Ingredients that Have Potential as Antidiabetic." *Proceeding of Mulawarman Pharmaceuticals Conferences*. Vol. 14. 2021.
- Nur'Afni Oktaviana Buhang, Siti Nuryanti, and Daud Karel Walanda. "UJI AKTIVITAS ANTIOKSIDAN EKSTRAK BAYAM MERAH (BLITUM RUBRUM) DALAM PELARUT ETANOL DAN AIR DENGAN PEREAKSI DPPH."
- Pasaribu, Fidayani, Panal Sitorus, and Saiful Bahri. "Uji ekstrak etanol kulit buah manggis (*Garcinia mangostana* L.) terhadap penurunan kadar glukosa darah." *Journal of Pharmaceutics and Pharmacology* 1.1 (2012): 1-8.
- Pop-Busui, Rodica, et al. "Heart failure: an underappreciated complication of diabetes. A consensus report of the American Diabetes Association." *Diabetes Care* 45.7 (2022): 1670-1690.
- Port, M.C. "Pathofiology Concep Of Altered Healt Status Second Edition". Lipincot Company. Phyladelpia. 1995:120-123
- Prisdiany, Yulin, et al. "Potensi Tanaman Herbal Antidiabetes untuk Minuman Obat: Sebuah Literatur Review." *Indonesian Journal of Clinical Pharmacy* 10.2 (2021): 144-158.
- Resti, Intan Apri, and Hesty Parbuntari. "Identifikasi Senyawa Metabolit Sekunder Ekstrak Jamur Tiram Putih (*Pleurotus ostreatus* L.)." *Periodic* 11.2 (2022): 65-69.
- Rinawati, Rinawati, E. Suharyanto, and Nastiti Wijayanti. "Pengaruh Ekstrak Rebusan Daun *Tithonia diversifolia* (Hemsl.) A. Gray Terhadap Kadar Glukosa Darah." *BIOTIK: Jurnal Ilmiah Biologi Teknologi dan Kependidikan* 7.1 (2019): 41-48.
- Rinawati, Rinawati, Sri Wahyuni Muhsin, and Siti Maisyaroh Fitri Siregar. "Pengaruh Ekstrak Air Selada Laut (*Ulva Lactuca*) Terhadap Berat Badan Pada Tikus Diabetes." *STIGMA: Jurnal Matematika dan Ilmu Pengetahuan Alam Unipa* 13.01 (2020): 39-46.
- Saeedi, Pouya, et al. "Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas." *Diabetes research and clinical practice* 157 (2019): 107843.
- Setiawan, B dan Suhartono, E. 2005. *Stres Oksidatif dan Peran Antioksidan pada Diabetes Mellitus*. Majalah Kedokteran Indonesia. 55(2):87
- Suryandari, Mercyska, and Galuh Gondo Kusumo. "Identifikasi Senyawa metabolit Sekunder Ekstrak Kulit Bawang Merah (*Allium cepa* L.) dari Berbagai Macam Pelarut." *Journal Pharmasci (Journal of Pharmacy and Science)* 7.2 (2022).
- Suryani, Nany. "Pengaruh Ekstrak Metanol Biji Mahoni terhadap Peningkatan Kadar Insulin, Penurunan Ekspresi TNF- $\alpha$  dan Perbaikan Jaringan Pankreas Tikus Diabetes." *Jurnal kedokteran brawijaya* 27.3 (2013): 137-145.

- Tair, Z. I., F. Bensalah, and F. Boukortt. "Effect of green alga *Ulva lactuca* polysaccharides supplementation on blood pressure and on atherogenic risk factors, in rats fed a high fat diet." *Annales de cardiologie et d'angiologie*. Vol. 67. No. 3. 2018
- Tandi, Joni, Tien Wahyu Handayani, and Agustinus Widodo. "Qualitative And Quantitative Determination Of Secondary Metabolites And Antidiabetic Potential Of *Ocimum Basilicum* L. Leaves Extract." *RASĀYAN J. Chem* 14.1 (2021): 622-628.
- Thongsom, Montakarn, et al. "Antioxidant and hypoglycemic effects of *Tithonia diversifolia* aqueous leaves extract in alloxan-induced diabetic mice." *Advances in Environmental Biology* (2013): 2116-2126.
- Touyz, Rhian M., et al. "Molecular Mechanisms Underlying Vascular Disease in Diabetes." *Blood Pressure Disorders in Diabetes Mellitus*. Springer, Cham, 2023. 105-118.
- Whiting, David R., et al. "IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030." *Diabetes research and clinical practice* 94.3 (2011): 311-321.
- WHO. 2018. Prevention of Diabetes Mellitus. Technical Report Series. Geneva.
- Zahra, Hafizh, et al. "Aktivitas Antiulseratif Berbagai Tanaman Herbal dan Prospek Masa Depan Sebagai Tanaman Budidaya: Anti-ulceritis Activity of Various Herbal Plants and Future Prospects as Cultivated Plants." *Jurnal Sains dan Kesehatan (J. Sains Kes.)* 4.3 (2022): 343-353.
- Zulfadhli, Zulfadhli, and Rinawati Rinawati. "Potensi Selada Laut *Ulva lactuca* Sebagai Antifungi Dalam Pengendalian Infeksi *Saprolegnia* Dan *Achlya* Pada Budidaya Ikan Kerling (*Tor sp.*)" *Jurnal Perikanan Tropis* 5.2 (2018): 183-188.

\*\*\*\*\*